

**Table A1. Healthy People 2000 Sexually Transmitted Diseases Objective 19.1–19.8 Status**

| <b>Objective</b> |  | <b>Baseline Year</b> | <b>Baseline</b>    | <b>1995</b>         | <b>1996</b> | <b>1997</b> | <b>1998</b> | <b>1999</b> | <b>2000</b> |
|------------------|--|----------------------|--------------------|---------------------|-------------|-------------|-------------|-------------|-------------|
| <b>19.1</b>      | <b>Gonorrhea (per 100,000 persons)</b>                       | 1989                 | 300                | 149                 | 124         | 123         | 133         | 133         | 100         |
| a.               | Black (non-Hispanic)   | 1989                 | 1,990              | 1,046               | 817         | 802         | 851         | 849         | 650         |
| b.               | Adolescents 15-19 years                                      | 1989                 | 1,123              | 671                 | 544         | 522         | 547         | 534         | 375         |
| c.               | Female 15-44 years   | 1989                 | 501                | 299                 | 259         | 252         | 282         | 283         | 175         |
| <b>19.2</b>      | <b>Chlamydia prevalence among females 15-24 years</b>        |                      |                    |                     |             |             |             |             |             |
|                  | Female 15-19 years   | 1988                 | 12.2%              | 6.7%                | 5.4%        | —           | 6.9%*       | 6.6%*       | 5%          |
|                  | Female 20-24 years   | 1988                 | 8.5%               | 4.2%                | 3.4%        | —           | 4.4%*       | 4.5%*       | 5%          |
| <b>19.3</b>      | <b>Primary and secondary syphilis (per 100,000 persons)</b>  | 1989                 | 18.1               | 6.3                 | 4.3         | 3.2         | 2.6         | 2.5         | 4           |
| a.               | Black  | 1989                 | 118                | 45                  | 30          | 22          | 17          | 15          | 30          |
| <b>19.4</b>      | <b>Congenital syphilis (per 100,000 live births)</b>         | 1990                 | 91.0               | 47.4                | 33.3        | 27.7        | 21.6        | 14.3        | 40          |
| a.               | Black  | 1992                 | <sup>a</sup> 417.8 | 213.2               | 150.5       | 122.4       | 90.3        | 57.9        | 175         |
| b.               | Hispanic   | 1992                 | <sup>a</sup> 134.6 | 61.2                | 38.9        | 33.4        | 28.7        | 20.4        | 50          |
| <b>19.5</b>      | <b>Annual number of first time consultations<sup>1</sup></b> |                      |                    |                     |             |             |             |             |             |
|                  | Genital herpes   | 1988                 | 163,000            | 160,000             | 208,000     | 176,000     | 188,000     | 224,000     | 138,500     |
|                  | Genital warts  | 1988                 | 290,000            | 253,000             | 191,000     | 145,000     | 211,000     | 240,000     | 246,500     |
| <b>19.6</b>      | <b>Pelvic inflammatory disease</b>                           |                      |                    |                     |             |             |             |             |             |
|                  | Hospitalizations per 100,000 females 15-44 years             | 1988                 | 311                | 162                 | 164         | 157         | 155         | —           | 100         |
|                  | Initial visits to physicians (number of visits) <sup>1</sup> | 1988                 | 430,800            | 262,000             | 286,000     | 261,000     | 234,000     | 251,000     | 290,000     |
|                  | Hospitalizations per 100,000 females                         |                      |                    |                     |             |             |             |             |             |
| a.               | Black 15-44 years  | 1988                 | 655                | 296                 | 320         | 281         | 291         | —           | 150         |
| b.               | Adolescents 15-19 years                                      | 1988                 | 342                | 141                 | 168         | 186         | 162         | —           | 110         |
| <b>19.7**</b>    | <b>Sexually transmitted Hepatitis B (number of cases)</b>    | 1987                 | 47,593             | <sup>2</sup> 29,446 | —           | —           | —           | —           | 30,500      |
| <b>19.8</b>      | <b>Repeat gonorrhea infection in last 12 months</b>          | 1987                 | 20%                | 18.4%               | 18.5%       | 17.0%       | 17.5%       | 17.2%       | 15%         |
| a.               | Black  | 1992                 | 21.3%              | 20.1%               | 19.8%       | 18.3%       | 18.6%       | 19.2%       | 17%         |

—Data not available.

<sup>a</sup>Baseline has been revised.

<sup>1</sup>As measured by first-time visits to physicians' offices.

<sup>2</sup>Data are provisional.

NOTE: Data include revisions and, therefore, may differ from data previously published in these reports and other publications.

## Data Sources

| Objective number | Data Source   |
|------------------|---|
| 19.1,19a-c       | Sexually Transmitted Disease Surveillance System, CDC, NCHSTP.  |
| 19.2             | Sexually Transmitted Disease Surveillance System, CDC, NCHSTP.  |
| 19.3, 19.3a      | Sexually Transmitted Disease Surveillance System, CDC, NCHSTP.  |
| 19.4             | Sexually Transmitted Disease Surveillance System, CDC, NCHSTP.  |
| 19.5             | National Disease and Therapeutic Index, IMS America, Ltd.   |
| 19.6, 19.6a-b    | For hospitalizations, National Hospital Discharge Survey, CDC, NCHS.<br>For number of visits, National Disease and Therapeutic Index, IMS America, Ltd. |
| 19.7**           | Viral Hepatitis Surveillance System, CDC, NCID.   |
| 19.8             | Gonococcal Isolate Surveillance Project, CDC, NCHSTP.   |

\*Positivity not adjusted for changes in laboratory test method in 1998-1999 and associated increases in test sensitivity.

\*\*Duplicate Objective.